

A SPITZER SEARCH FOR SUBSTELLAR COMPANIONS TO LOW MASS WHITE DWARFS*

MUKREMIN KILIC^{1,2}, WARREN R. BROWN, AND B. MCLEOD
 Smithsonian Astrophysical Observatory, 60 Garden St., Cambridge, MA 02138
 Draft version November 4, 2009

ABSTRACT

The formation scenarios for *single* low-mass ($M < 0.45M_{\odot}$) white dwarfs include enhanced mass loss from a metal-rich progenitor star or a common envelope phase of a solar-like star with a close-in massive planet or a brown dwarf. Both scenarios suggest that low-mass white dwarfs may have planets. Here, we present a *Spitzer* IRAC search for substellar and planetary mass companions to 14 low-mass white dwarfs. One of our targets, HS 1653+7753, displays near- and mid-infrared flux excess. However, follow-up MMT observations show that this excess is due to a nearby resolved source, which is mostly likely a background object. Another target, PG 2257+162, shows flux excess compatible with a late-type stellar companion. We do not detect substellar companions to any of the remaining targets. In addition, eight of these stars do not show any radial velocity variations, ruling out stellar mass companions including other white dwarfs. We conclude that a significant fraction of the low-mass white dwarfs in our sample do not have stellar or massive brown dwarf companions.

Subject headings: infrared: stars — planetary systems — stars: low-mass, brown dwarfs — white dwarfs

1. INTRODUCTION

The fate of planetary systems around Sun-like stars can be studied by finding planets around their white dwarf (WD) remnants. WDs are physically small ($R \sim 0.01R_{\odot}$), which sharply limits their luminosity in the infrared, where the planet spectrum peaks (Burleigh et al. 2002; Ignace 2001). Compared to main-sequence (MS) stars, the significant gain in contrast makes WDs excellent targets for photometric searches for planetary companions. However, to this day, no planet detection around WDs has been reported. *Spitzer* IRAC imaging surveys by Mullally et al. (2007), Debes et al. (2007), Farihi et al. (2008), and Kilic et al. (2009b), and near-infrared imaging surveys by Friedrich et al. (2006), Burleigh et al. (2008), and Hogan et al. (2009) were unsuccessful in finding planetary mass companions to WDs with limits typically between 5 and $10M_J$. Here we focus on using low-mass WDs to provide additional leverage for detecting planetary mass companions.

1.1. CURRENT CONSTRAINTS

Timing measurements of pulsating WDs is an alternative, powerful method to search for planets (Winget et al. 2003), but this method can be used only for a limited sample of stars. There is a promising signature of a $M \sin i = 2.4M_J$ planet in the pulsation timing measurements of the WD GD 66 (Mullally et al. 2009). However, a full orbit has not been observed, and this detection remains provisional (F. Mullally 2009, priv. comm.).

The discovery of circumstellar debris disks around more than a dozen WDs imply that 1% to 3% of WDs with cooling ages < 500 Myr have disks (see Farihi et al. 2009, and references therein). These disks most likely formed by tidally disrupted asteroids (Jura 2003). Hence, there is indirect evidence that at least a few percent of WDs have remnant planetary systems. Of course, the observed debris disk fraction

may represent only the planetary systems that become unstable during post-MS evolution, and the true fraction of WDs with planets may be much higher.

The reason for the lack of known planetary companions to WDs may be a target selection effect (Kilic et al. 2007b). Fischer & Valenti (2005) find that less than 3% of the stars with $-0.5 < [\text{Fe}/\text{H}] < 0.0$ have Doppler-detected planets, whereas 25% of the stars with $[\text{Fe}/\text{H}] > +0.3$ have gas giant planets. Based on the age-metallicity relation derived by Reid et al. (2007), Kilic et al. (2007b) find that only 3% of the stars in the solar neighborhood have had $[\text{Fe}/\text{H}] > +0.3$ in the past 10 Gyr. Therefore, a random selection of WDs would be dominated by WDs that come from low metallicity progenitors. The best way to find planetary companions to WDs may be to search for companions to WDs that form from metal-rich stars.

1.2. LOW-MASS WHITE DWARFS

Low-mass helium-core WDs ($M < 0.45 M_{\odot}$) can be produced from interacting binary systems, and traditionally all of them have been attributed to this channel. However, a low-mass WD can also result from a single star that experiences severe mass loss on the first ascent giant branch. Kalirai et al. (2007) identified several low-mass WDs in the upper part of the WD cooling sequence of the metal-rich cluster NGC 6791, and they argued that there is a mechanism in clusters to produce low-mass WDs without requiring binary star interactions. However, this argument is now challenged by Bedin et al. (2008), who argue that the bright peak observed in the WD luminosity function of this cluster is better explained as binary WDs. Bedin et al. (2008) also argue that the low-mass WDs found by Kalirai et al. (2007) can be explained as the product of binary star evolution.

Unlike the relatively faint low-mass WDs in NGC 6791, it is possible to obtain radial velocity measurements of low-mass WDs in the field and test the binary formation scenario. Around half the low-mass WDs observed by Maxted et al. (2000) lack detectable radial velocity companions. The ESO SNIa Progenitor Survey (SPY) searched for radial velocity variations in more than a thousand WDs using the Very Large Telescope (Napiwotzki et al. 2001). They found that about

*BASED PARTLY ON OBSERVATIONS OBTAINED AT THE MMT OBSERVATORY, A JOINT FACILITY OF THE SMITHSONIAN INSTITUTION AND THE UNIVERSITY OF ARIZONA.

¹ *Spitzer* Fellow

² mkilic@cfa.harvard.edu

58% of the low-mass WDs in their sample do not show radial velocity variations (Napiwotzki et al. 2007). Their detailed analysis of the detection probability of MS, WD, and brown dwarf companions show that their observations of low-mass WDs are inconsistent with the idea that all of them reside in close binary systems. The SPY observations are less sensitive to brown dwarf companions, because of their smaller mass, but even in this case 100% binary fraction rate can be ruled out (R. Napiwotzki 2007, private communication). Kilic et al. (2007b) argue that these single low-mass WDs might come from old metal-rich stars that truncate their evolution prior to the helium flash from severe mass loss. The fraction of single low-mass WDs among the field WD sample is also compatible with the fraction of metal-rich dwarfs among the field dwarf population.

A strong implication of the Kilic et al. (2007b) scenario is that single low-mass WDs are good targets for planet searches because they may arise from metal-rich progenitors. This leads to an inversion of prior strategies for seeking planetary companions to WDs, which avoid low-mass WDs on the grounds that they are the products of binary interactions. Instead, some of these objects may be the descendants of metal-rich stars that are known to have a high frequency of planet detection. If the progenitors of these single low-mass WDs have planets, the planets are more likely to survive the post-MS evolution, as these He-core WDs do not go through the asymptotic giant branch phase.

An alternative formation scenario for single low-mass WDs is proposed by Nelemans & Tauris (1998). In this scenario, low-mass WDs form from Sun-like stars with close planetary companions. A common envelope phase with a massive planet can expel the stellar envelope and form a low-mass WD. The planet may (not) survive the common envelope phase. However, if there are additional planets in wide orbits, they are likely to survive (Livio & Soker 1983; Burleigh et al. 2002; Duncan & Lissauer 1998).

Both scenarios proposed by Nelemans & Tauris (1998) and Kilic et al. (2007b) suggest that low-mass WDs may have planets. Here we present a *Spitzer* IRAC search for substellar companions to 14 low-mass WDs. Our sample selection and observations are discussed in Section 2, whereas the detectability of planets around our targets and the results of our search are discussed in Section 3 and 4.

2. OBSERVATIONS

Maxted et al. (2000) presented a radial velocity survey of 71 WDs including 13 low-mass WDs with no detectable radial velocity variations. Hansen et al. (2006) and Mullally et al. (2007) obtained IRAC observations of 3 of these stars (WD 0316+345, WD 1353+409, and WD 1614+136). We target the remaining 10 single low-mass WDs for follow-up observations. In addition, we target 4 low-mass WDs from the Palomar-Green Survey (Liebert et al. 2005). Farihi et al. (2005) presented near-infrared photometry for two of these stars which did not reveal any companions. The physical parameters of our targets, including masses, distances, and WD cooling ages, are presented in Table 1. All of our targets are hydrogen-rich DA WDs.

Observations reported here were obtained as part of the Cycle 5 *Spitzer* Fellowship Program 474. We obtained 3.6, 4.5, 5.8, and 8 μm images with integration times of 30 or 100 seconds per dither, with five or nine dithers per object. We use the IRAF PHOT and IDL astrolib packages to perform aperture photometry on the individual BCD frames from the latest

TABLE 1
LOW MASS WD TARGETS

Object	T_{eff} (K)	M_{WD} ($M_{\odot}$)	d (pc)	τ_{WD} (Myr)	Reference
PG 0132+254	19960	0.41	180	60	1
PG 0237+242	69160	0.40	881	20	1
WD 0339+523	12640	0.33	103	180	2
WD 0437+152	10410	0.38	153	340	3
WD 0453+418	13660	0.44	44	150	2
WD 0509-007	32200	0.40	117	10	4
PG 0808+595	27330	0.42	238	10	1
PG 0943+441	12820	0.41	30	340	1
WD 0950-572	12400	0.44	58	230	2
PG 1229-013	19430	0.41	59	70	1
HS 1653+7753	29400	0.32	180	10	5
PG 1654+637	15070	0.44	98	200	1
PG 2226+061	15280	0.44	65	190	1
PG 2257+162	24580	0.43	212	20	1

NOTE. — 1- Liebert et al. (2005), 2- Gianninas et al. (2005), 3- Moran (1999), 4- Vennes et al. (1997), 5- Homeier et al. (1998)

available IRAC reduction pipeline (17.2 or 18.5 for our targets). Since our targets are relatively faint, we use the smallest aperture (2 pixels) for which there are published aperture corrections. Following the IRAC calibration procedure, corrections for the location of the source in the array were taken into account before averaging the fluxes of each of the dithered frames at each wavelength. Channel 1 (3.6 μm) photometry is also corrected for the pixel-phase-dependence. The results from IRAF and IDL reductions are consistent within the errors. The photometric error bars are estimated from the observed scatter in the 5 (or 9) images corresponding to the dither positions. We also add the 3% absolute calibration error in quadrature. Finally, we divide the estimated fluxes by the color corrections for a Rayleigh-Jeans spectrum (Reach et al. 2005). These corrections are 1.0111, 1.0121, 1.0155, and 1.0337 for the 3.6, 4.5, 5.8, and 8 μm bands, respectively.

We present the IRAC photometry of our targets in Table 2. Most of our targets are detected in 2MASS, at least in the *J* band. Several targets also have Sloan Digital Sky Survey (SDSS) photometry available. We obtained *V* and *I* band photometry of HS 1653+7753 using the MDM 2.4m telescope equipped with the Echelle CCD on UT 2009 April 5. We use observations of the standard star field PG 1633+099 (Landolt 1992) to calibrate the photometry. The photometric reductions were performed by J. Thorstensen, and kindly made available to us. HS 1653+7753 has *V* = 15.08 mag and *I* = 15.33 mag. We estimate an internal accuracy of 0.01 mag for the photometry, but to be conservative, we adopt errors of 0.03 mag. H. C. Harris kindly provided us unpublished *BVI* photometry of WD 0339+523. We use the optical photometry from McCook & Sion (2006) for the remaining targets.

In addition, we obtained *JHK* imaging observations of HS 1653+7753 using the MMT and Magellan Infrared Spectrograph (MMIRS; McLeod et al. 2004) on the MMT on UT 2009 June 13. The observations were obtained under cloudy conditions, and they included 10 exposures of 30 s in *J*, 10 exposures of 20 s in *H*, and 25 exposures of 45 s in *K*.

3. RESULTS

Figure 1 presents the spectral energy distributions of twelve of our fourteen low-mass WDs. Two WDs, with more complicated SEDs, are presented below. D. Koester kindly pro-

TABLE 2
 IRAC PHOTOMETRY OF LOW MASS WDs

Object	3.6 μ m (μ Jy)	4.5 μ m (μ Jy)	5.8 μ m (μ Jy)	8.0 μ m (μ Jy)
PG 0132+254	54.7 $\pm$ 3.2	32.9 $\pm$ 1.4	21.1 $\pm$ 11.0	25.6 $\pm$ 11.2
PG 0237+242	54.2 $\pm$ 3.4	34.4 $\pm$ 1.8	24.2 $\pm$ 10.8	22.5 $\pm$ 8.4
WD 0339+523	103.9 $\pm$ 5.0	66.1 $\pm$ 3.8	33.7 $\pm$ 19.2	23.7 $\pm$ 20.6
WD 0437+152	79.0 $\pm$ 4.1	49.2 $\pm$ 2.2	31.3 $\pm$ 17.7	22.0 $\pm$ 19.3
WD 0453+418	449.7 $\pm$ 14.4	287.3 $\pm$ 10.4	176.7 $\pm$ 25.9	102.9 $\pm$ 46.8
WD 0509-007	333.3 $\pm$ 11.1	213.4 $\pm$ 11.5	141.8 $\pm$ 18.9	73.5 $\pm$ 18.3
PG 0808+595	48.3 $\pm$ 5.3	27.8 $\pm$ 2.4	13.5 $\pm$ 13.2	13.4 $\pm$ 4.2
PG 0943+441	881.2 $\pm$ 32.4	549.1 $\pm$ 19.4	367.1 $\pm$ 16.7	212.7 $\pm$ 19.7
WD 0950-572	269.6 $\pm$ 14.4	173.7 $\pm$ 7.7	121.9 $\pm$ 19.3	121.9 $\pm$ 40.1
PG 1229-013	237.6 $\pm$ 9.2	150.3 $\pm$ 6.8	80.4 $\pm$ 9.4	73.8 $\pm$ 32.1
HS 1653+7753	161.3 $\pm$ 5.1	106.5 $\pm$ 3.9	66.9 $\pm$ 5.7	39.5 $\pm$ 7.9
PG 1654+637	89.6 $\pm$ 4.7	53.9 $\pm$ 2.3	37.1 $\pm$ 14.6	15.2 $\pm$ 11.1
PG 2226+061	198.6 $\pm$ 14.6	122.1 $\pm$ 7.1	74.5 $\pm$ 24.6	41.2 $\pm$ 21.0
PG 2257+162	410.7 $\pm$ 17.9	282.4 $\pm$ 14.1	190.1 $\pm$ 30.4	113.2 $\pm$ 36.6

vided pure-hydrogen atmosphere WD model spectra for our targets. We perform synthetic photometry on these models using the appropriate transmission curves for the *BVRI*, SDSS, 2MASS, and IRAC filters. The resulting fluxes are then compared with the observations to find the normalization factor for the WD models. We weight fluxes by their associated error bars. The solid lines in Figure 1 present the appropriate WD model for each star normalized to match the observations. The use of optical and near-infrared photometry helps us constrain the predicted mid-infrared photospheric fluxes for WDs. The mid-infrared photometry of all but one of the twelve WDs presented in this figure is consistent with the predicted photospheric flux from WDs; we do not detect excess mid-infrared flux from planets or substellar companions. The thermal emission from planetary companions would be detected most strongly at 4.5 μ m (Burrows et al. 2003).

One of the objects in Figure 1, WD 0950-572 shows a mid-infrared flux excess. Figure 2 presents a contour map of the immediate field around this WD; it is near a bright star and even the smallest aperture that we use (2 pixels) is contaminated by flux from this nearby bright star. Therefore, the observed excess flux is most likely not intrinsic to the WD environment.

3.1. DETECTABILITY OF PLANETS

The usual method for estimating the upper mass limit for substellar companions in photometric surveys is to calculate the total age of the white dwarf, i.e. the WD cooling age plus the MS lifetime, and then use evolutionary models for substellar objects and giant planets to predict their brightness at the age and distance of the WD (see for example Hogan et al. 2009; Farihi et al. 2008). But there is a problem for low mass WDs: if they form through common envelope evolution, it is not possible to know their progenitor mass and their MS lifetimes. Hence only a lower limit to the age of the system, i.e. the WD cooling age, is known. Even if the low-mass WDs form directly from single metal-rich progenitors, currently we do not have initial-final mass relations that would give us progenitor masses and lifetimes for such systems. Therefore, all we can do is to calculate companion upper mass limits for a range of possible ages starting from the WD cooling ages to several Gyr. All of the objects in our sample are hotter than 10,000 K (Table 1), therefore their WD cooling ages are less than 350 Myr (Bergeron et al. 1995).

A relatively massive planet or brown dwarf can stay hot




FIG. 1.— Spectral energy distributions of 12 low-mass WDs. The expected photospheric flux from the WDs are shown as solid lines (D. Koester 2009, priv. comm.).




FIG. 2.— Contour map of the immediate field around WD 0950–572. The map covers 20 x 20 pixels, which corresponds to 24" x 24". Dashed circle marks a 2 pixel aperture centered on the WD.

and easily detectable by *Spitzer* for billions of years. Some brown dwarfs or planets may also be reheated during the red giant phase of the progenitor’s evolution, either by direct radiation or by impacts from planets and debris belts destabilized during stellar mass loss (Parriott & Alcock 1998; Debes & Sigurdsson 2002). Collisions between two planets may create mergers that would essentially reset the cooling clock of the planet and make the remnant planet overluminous by a factor of a hundred. However, Debes & Sigurdsson (2002) predict that only about 2% of young WDs may have overluminous planetary companions that form through collisions.

Melis et al. (2009) report the discovery of a unique first-




FIG. 3.— Spectral energy distribution of PG 2226+061 compared to the expected photospheric flux from the WD (solid line), and from planetary (1–25 M_J) and brown dwarf companions (dotted lines, Burrows et al. 2003, 2006). The red cross marks the 3σ upper limit of the 4.5 μm photometry.

ascent giant star with a substantial dust disk. They suggest that the circumstellar material around this star may originate from a common-envelope phase with a low-mass stellar or substellar companion. The gaseous and dusty material is similar to the primordial disks observed around T Tauri stars. They speculate that these disks may form a second generation of planets around first-ascent giant stars. Single low-mass WDs may descend from such stars (Nelemans & Tauris 1998). The discovery of a close-in substellar companion to the low-mass WD 0137–049 implies that this substellar companion has gone through a common-envelope phase that resulted in the ejection of the envelope of the progenitor first-ascent giant star (Maxted et al. 2006). If the first-ascent giant stars with gaseous/dusty disks are common and if these stars form low-mass single WDs, then these WDs may have second generation planets as well. These planets would be as young as the cooling age of the WD, and therefore warmer and more luminous.

3.2. LIMITS ON COMPANIONS

For the 11 stars presented in Figure 1 (excluding WD 0950–572), we use Baraffe et al. (2003) and Burrows et al. (2003, 2006) models to put upper limits on companions that would escape detection. I. Baraffe kindly provided us substellar cooling models updated to include fluxes in the IRAC bandpasses for 0.1–5 Gyr old objects with $M = 1\text{--}100 M_J$.

Figure 3 shows an example of predicted excess flux from planetary and brown dwarf companions around one of our targets, PG 2226+061. By demanding a 3σ excess from any possible companion, we exclude 100 Myr old, $M > 4M_J$ planets around PG 2226+061. Depending on the unknown ages of the possible planetary companions, this limit goes up to more than 30 M_J for 5 Gyr or older planets. In all cases, we are able to rule out brown dwarf companions with $T_{\text{eff}} \geq 700$ K around this WD.

We achieve similar limits on substellar companions for the rest of our sample. These limits and our search radii are presented in Table 3. Since the total ages (WD cooling age + MS lifetime) are unknown, we present companion mass limits for a range of ages. If the planets are as old as the WD cooling ages (20–340 Myr), we can rule out planets more massive than 5 M_J to 10 M_J around 10 of our targets; second-generation massive planets are not common around low-mass WDs. However, if the planets are as old as 5 Gyr, only massive brown dwarf companions are ruled out.

3.3. HS 1653+7753

One of our targets, HS 1653+7753, shows excess flux in the mid-infrared. The contour maps of the IRAC images show that there is a nearby faint source in the IRAC beam. The 2MASS images do not reveal any nearby, faint sources. We obtained MMIRS *JHK* imaging of HS 1653+7753 to resolve this nearby source. Figure 4 displays *JHK* and IRAC 4.5 μm images of the immediate field around HS 1653+7753. Our MMIRS data shows that there is a faint source to the southeast of the WD at 2.2" separation. This source is relatively bright in the mid-infrared, and contaminates the IRAC photometry even in a 2 pixel aperture. We use a 3 pixel (3.6") aperture to measure the total flux from the two sources in our IRAC data.

The nearby source is clearly resolved in the *J*-band data, but it is not detected well in the *H* and *K* band images. This prevents us from using point spread function fitting photometry. The FWHM of the near-infrared images is 1.2" (6 pixels). We

TABLE 3
 RULED OUT COMPANIONS

Object	R (AU)	0.1 Gyr (M_J)	0.5 Gyr (M_J)	1.0 Gyr (M_J)	5.0 Gyr (M_J)	BD ($T_{\text{eff}} \geq$)
PG 0132+254	430	> 5	> 10	> 15	> 40	800 K
PG 0237+242	2110	> 50	> 90	> 100	> 100	...
WD 0339+523	250	> 5	> 10	> 15	> 40	800 K
WD 0437+152	370	> 6	> 10	> 20	> 40	800 K
WD 0453+418	110	> 4	> 9	> 12	> 30	700 K
WD 0509-007	280	> 10	> 40	> 50	> 72	1600 K
PG 0808+595	570	> 9	> 20	> 30	> 60	1100 K
PG 0943+441	70	> 2	> 7	> 10	> 20	700 K
PG 1229-013	140	> 2	> 6	> 9	> 20	700 K
PG 1654+637	240	> 2	> 5	> 7	> 15	700 K
PG 2226+061	160	> 4	> 10	> 15	> 30	700 K

NOTE. — The mass limits are based on the models of I. Baraffe (2009, private communication; Baraffe et al. 2003). The temperature limits for brown dwarf companions (the last column) are based on the models of Burrows et al. (2006).

use a $1.2''$ aperture to measure the flux from the WD only, and a $3.6''$ aperture to measure the total flux from the two sources. The $6.8' \times 6.8'$ field of view of MMIRS enables us to use about 20 nearby 2MASS stars to calibrate the photometry. We checked the photometry for several 2MASS stars using $1.2''$ and $3.6''$ apertures, and found them to be consistent within the errors. We measure $J = 15.80 \pm 0.01$, $H = 15.91 \pm 0.02$, and $K = 16.06 \pm 0.03$ mag for the WD (using a $1.2''$ aperture), and $J = 15.70 \pm 0.02$, $H = 15.71 \pm 0.05$, and $K = 15.83 \pm 0.07$ mag for the total flux in a $3.6''$ aperture. The latter are consistent with 2MASS photometry for HS 1653+7753 within the errors.

Figure 5 presents the spectral energy distribution of HS 1653+7753. We show the near-infrared photometry for both $1.2''$ and $3.6''$ apertures, and the mid-infrared photometry for a $3.6''$ aperture. The combined flux from the WD and possible brown dwarf companions (Burrows et al. 2006) are shown in the top panel. If the two objects are bound, the observed mid-infrared flux excess implies a relatively warm brown dwarf. However, the excess flux (at $4.5 \mu\text{m}$) is not explained well by




FIG. 4.— MMIRS J, H, K-band and IRAC $4.5 \mu\text{m}$ (top left to bottom right) images of the immediate field around HS 1653+7753. North is up and East is to the left. The image covers $28'' \times 28''$. The circle marks a $3.6''$ aperture centered on the WD.




FIG. 5.— Spectral energy distribution of HS 1653+7753. The near-infrared photometry using both $1.2''$ (lower points) and $3.6''$ (upper points) apertures is shown. The solid lines show the expected photospheric flux from the WD. The dashed lines show the combined flux from the WD and 1600 K - 2000 K brown dwarf companions (top panel), and a background object with a blackbody temperature of 2200 K (bottom panel).

these models. The bottom panel in Figure 5 presents the combined flux from the WD and a background object with a blackbody temperature of 2200 K. The mid-infrared flux excess is explained better by a 2200 K background object. Therefore, the resolved object near the WD is most likely a background source.

HS 1653+7753 has $\mu_{\text{RA}} = -108 \pm 32$ and $\mu_{\text{DEC}} = 118 \pm 99$ mas yr $^{-1}$ in the USNO-B catalog (Monet et al. 2003). Using 2MASS and MMIRS J -band images, we measure $\mu_{\text{RA}} = -19 \pm 15$ and $\mu_{\text{DEC}} = 76 \pm 19$ mas yr $^{-1}$. If HS 1653+7753 and the nearby source are not bound, they will be moving further apart. Follow up J -band observations in a few years will show if this system is bound.

3.4. PG 2257+162

PG 2257+162 is classified as a tentative WD + low-mass MS binary by Wachter et al. (2003) based on its 2MASS near-infrared colors. It shows significant flux excesses in the near- and mid-infrared, and it is not resolved in the IRAC images. The spectral energy distribution of this WD is presented in Figure 6. It is clear from this figure that the excess comes from an object with a blackbody temperature of 3300 K. If the unresolved companion is physically associated with the WD, it is most likely a low-mass stellar companion. We are currently obtaining radial velocity observations of this WD using the 1.5m Tillinghast telescope to see if this system is a physical binary or not.

4. DISCUSSION AND CONCLUSIONS

Our IRAC survey of 14 low-mass WDs did not reveal any substellar or planetary mass companions. One of our targets, HS 1653+7753, has a nearby source that is likely to be a background object. Another target, PG 2257+162, possibly has a




FIG. 6.— Spectral energy distribution of PG 2257+162. The star is unresolved in the IRAC images. The excess infrared flux is likely due to a late type star companion. The solid, dashed, and dotted lines show the expected flux for the WD, a 3300 K blackbody, and the combination of the two, respectively.

low-mass stellar companion. In addition, the IRAC photometry for one of our targets, WD 0950–572 is affected by a nearby star. The remaining 11 low-mass WDs do not show any mid-infrared flux excesses. Adding the three low-mass WDs observed by Hansen et al. (2006) and Mullally et al. (2007) to our sample brings the sample size to 14 WDs with no detectable mid-infrared flux excess. Eleven of these targets do not show radial velocity variations, ruling out companions more massive than $\approx 0.08 M_{\odot}$ (Maxted et al. 2000). Therefore, these 11 low-mass WDs appear to have no stellar (including another WD) or substellar companions to the limits given in Table 3. The non-detection of stellar, substellar, or planetary companions to our targets is not inconsistent with the idea that single low-mass WDs descend from single metal-rich stars through extreme mass loss.

Nelemans & Tauris (1998) predict that if a substellar object with $M < 32 M_J$ goes through a common-envelope phase with a star, it will either evaporate or spiral-in during the process. More massive objects are expected to survive this evolution, as evidenced by WD 0137–049B ($0.053 M_{\odot}$, Maxted et al. 2006), although the theoretical $32 M_J$ limit needs to be tested by observations. Nelemans & Tauris (1998) use WD 1614+136 to demonstrate that their common envelope phase scenario is feasible. They assume that WD 1614+136 is a remnant of a $1 M_{\odot}$ star with a core-mass of $0.33 M_{\odot}$. They

find that the minimum companion mass required to expel the envelope of the progenitor is $21 M_J$. No companion is detected in IRAC observations of this star, and Hansen et al. (2006) rule out $M \geq 20 M_J$ companions for ages ≤ 1 Gyr. This limit goes up to about $40 M_J$ for 5 Gyr old companions. A well tuned scenario involving a common envelope phase between a ~ 5 Gyr old $1 M_{\odot}$ star and a 21 – $40 M_J$ brown dwarf at a certain orbital separation can explain the single, low-mass WD 1614+136.

Our survey is sensitive to massive brown dwarf companions around the majority of our targets even at 5 Gyr (see Table 3). We rule out $M > 40 M_J$ companions around 8 of our targets at 5 Gyr. Like WD 1614+136, only a well tuned common-envelope phase scenario involving 20 – $40 M_J$ companions would explain the low mass WDs in our sample. Grether & Lineweaver (2006) find that the driest part (the minimum number) of the brown dwarf desert occurs at $31 M_J$. Thus, it seems unlikely that the scenario proposed by Nelemans & Tauris (1998) will explain all single, low-mass WDs. However, given the caveats in understanding the common-envelope phase evolution and our age dependent limits on possible companions, we cannot rule out this scenario.

An alternative scenario for the formation of single low-mass WDs involves binary mergers of even lower mass WDs (Iben et al. 1997). Maxted & Marsh (1998) find that such mergers result in rotational velocities on the order of 1000 km s^{-1} , unless the WDs are efficient in losing angular momentum through mass loss during the merging process. Maxted & Marsh (1998) measure rotational velocities of 50 km s^{-1} for two single low-mass WDs and they suggest that efficient angular momentum loss should take place if these WDs formed through binary mergers. Nelemans & Tauris (1998) find this formation scenario unlikely. Currently, there are no known binary low-mass WD progenitor systems with combined mass $\approx 0.4 M_{\odot}$ and with merger time shorter than a Hubble time (see Table 1 in Nelemans & Tout 2005). Radial velocity studies of extremely low-mass ($0.2 M_{\odot}$) WDs are therefore important to search for such systems (Kilic et al. 2007a, 2009a) and to see if this scenario is likely to explain at least a fraction of single, low-mass WDs.

Support for this work was provided by NASA through the Spitzer Space Telescope Fellowship Program, under an award from Caltech. We thank the MMIRS commissioning team for obtaining the observations, and T. von Hippel for a careful reading of an earlier version of this manuscript. We also thank an anonymous referee for useful suggestions.

REFERENCES

- Baraffe, I., Chabrier, G., Barman, T. S., Allard, F., & Hauschildt, P. H. 2003, *A&A*, 402, 701
- Bedin, L. R., Salaris, M., Piotto, G., Cassisi, S., Milone, A. P., Anderson, J., & King, I. R. 2008, *ApJ*, 679, L29
- Bergeron, P., Wesemael, F., & Beauchamp, A. 1995, *PASP*, 107, 1047
- Burleigh, M. R., Clarke, F. J., & Hodgkin, S. T. 2002, *MNRAS*, 331, L41
- Burleigh, M. R., et al. 2008, *MNRAS*, 386, L5
- Burrows, A., Sudarsky, D., & Lunine, J. I. 2003, *ApJ*, 596, 587
- Burrows, A., Sudarsky, D., & Hubeny, I. 2006, *ApJ*, 640, 1063
- Debes, J. H., & Sigurdsson, S. 2002, *ApJ*, 572, 556
- Debes, J. H., Sigurdsson, S., & Hansen, B. 2007, *AJ*, 134, 1662
- Duncan, M. J., & Lissauer, J. J. 1998, *Icarus*, 134, 303
- Farihi, J., Becklin, E. E., & Zuckerman, B. 2005, *ApJS*, 161, 394
- Farihi, J., Becklin, E. E., & Zuckerman, B. 2008, *ApJ*, 681, 1470
- Farihi, J., Jura, M., & Zuckerman, B. 2009, *ApJ*, 694, 805
- Fischer, D. A. & Valenti, J. 2005, *ApJ*, 622, 1102
- Friedrich, S., Zinnecker, H., Correia, S., Brandner, W., Burleigh, M., & McCaughrean, M. 2006, *ASP Conference Series*, 999
- Grether, D., & Lineweaver, C. H. 2006, *ApJ*, 640, 1051
- Hansen, B. M. S., Kulkarni, S., & Wiktorowicz, S. 2006, *AJ*, 131, 1106
- Hogan, E., Burleigh, M. R., & Clarke, F. J. 2009, *MNRAS*, in press
- Iben, I. J., Tutukov, A. V., & Yungelson, L. R. 1997, *ApJ*, 475, 291
- Ignace, R. 2001, *PASP*, 113, 1227
- Jura, M. 2003, *ApJ*, 584, L91
- Kalirai, J. S., Bergeron, P., Hansen, B. M. S., Kelson, D. D., Reitzel, D. B., Rich, R. M., & Richer, H. B. 2007, *ApJ*, 671, 748
- Kilic, M., Brown, W. R., Allende Prieto, C., Pinsonneault, M. H., & Kenyon, S. J. 2007a, *ApJ*, 664, 1088
- Kilic, M., Stanek, K. Z., & Pinsonneault, M. H. 2007b, *ApJ*, 671, 761

- Kilic, M., Brown, W. R., Allende Prieto, C., Swift, B., Kenyon, S. J., Liebert, J., & Agüeros, M. A. 2009a, *ApJ*, 695, L92
- Kilic, M., Gould, A., & Koester, D. 2009b, *ApJ*, 705, 1219
- Koester, D., Rollenhagen, K., Napiwotzki, R., Voss, B., Christlieb, N., Homeier, D., & Reimers, D. 2005, *A&A*, 432, 1025
- Landolt, A. U. 1992, *AJ*, 104, 340
- Lépine, S., & Shara, M. M. 2005, *AJ*, 129, 1483
- Liebert, J., Bergeron, P., & Holberg, J. B. 2005, *ApJS*, 156, 47
- Livio, M., & Soker, N. 1983, *A&A*, 125, L12
- Maxted, P. F. L., & Marsh, T. R. 1998, *MNRAS*, 296, L34
- Maxted, P. F. L., Marsh, T. R., & Moran, C. K. J. 2000, *MNRAS*, 319, 305
- Maxted, P. F. L., Napiwotzki, R., Dobbie, P. D., & Burleigh, M. R. 2006, *Nature*, 442, 543
- McCook, G. P., & Sion, E. M. 2006, *VizieR Online Data Catalog*, 3235, 0
- McLeod, B. A., Fabricant, D., Geary, J., Martini, P., Nystrom, G., Elston, R., Eikenberry, S. S., & Epps, H. 2004, *Proc. SPIE*, 5492, 1306
- Melis, C., Zuckerman, B., Song, I., Rhee, J. H., & Metchev, S. 2009, *ApJ*, 696, 1964
- Monet, D. G., et al. 2003, *AJ*, 125, 984
- Mullally, F., Kilic, M., Reach, W. T., Kuchner, M. J., von Hippel, T., Burrows, A., & Winget, D. E. 2007, *ApJS*, 171, 206
- Mullally, F., Reach, W. T., Degennaro, S., & Burrows, A. 2009, *ApJ*, 694, 327
- Napiwotzki, R., et al. 2001, *Astronomische Nachrichten*, 322, 411
- Napiwotzki, R., et al. 2007, 15th European Workshop on White Dwarfs, 372, 387
- Nelemans, G., & Tauris, T. M. 1998, *A&A*, 335, L85
- Nelemans, G., & Tout, C. A. 2005, *MNRAS*, 356, 753
- Parriott, J., & Alcock, C. 1998, *ApJ*, 501, 357
- Reach, W. T., et al. 2005, *PASP*, 117, 978
- Reid, I. N., Turner, E. L., Turnbull, M. C., Mountain, M., & Valenti, J. A. 2007, *ApJ*, 665, 767
- Wachter, S., Hoard, D. W., Hansen, K. H., Wilcox, R. E., Taylor, H. M., & Finkelstein, S. L. 2003, *ApJ*, 586, 1356
- Winget, D. E., et al. 2003, *Scientific Frontiers in Research on Extrasolar Planets*, 294, 59